




SÉRIE TRIGONOMÉTRICA
DE FOURIER ∞



$$S'G = F_1(x, y) + F_2(x, y)$$




$$P(x, y, \alpha)$$



$$P(x, y)$$

SISTEMA
CARTESIANO



$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos\left(\frac{n\pi}{L}x\right) + b_n \sin\left(\frac{n\pi}{L}x\right) \right)$$

$-L \leq x \leq L$

STF. $a_0 = \frac{1}{L} \int_{-L}^L f(x) dx$

$$a_n = \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{n\pi}{L}x\right) dx$$

$$b_n = \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{n\pi}{L}x\right) dx$$




Uma $f(t)$ $-L \leq t \leq L$ será PAR

$$f(-t) = f(t)$$

uma $f(t)$ $-L \leq t \leq L$ será IMPAR


PAR $f(-t) = -f(t)$

$$\int_{-L}^L f dt = 2 \int_0^L f dt$$




IMPAR


$$\int_{-L}^L f dt = 0$$



PAR



IMPAR



$$\langle \text{PAR} \rangle \langle \text{PAR} \rangle = \langle \text{PAR} \rangle$$

$$\langle \text{IMPAR} \rangle \langle \text{IMPAR} \rangle = \langle \text{PAR} \rangle$$

$$\langle \text{PAR} \rangle \langle \text{IMPAR} \rangle = \langle \text{IMPAR} \rangle$$

$$\int_{-L}^L \text{IMPAR} = 0$$

$$\int_{-L}^L \text{PAR} = 2 \int_0^L \text{PAR}$$

