

> restart

PROBLEMA DE LA CUERDA DE GUITARRA

> $EDP := \text{diff}(y(x, t), t\$2) - c \cdot 2 \cdot \text{diff}(y(x, t), x\$2) = 0$

$$EDP := \frac{\partial^2}{\partial t^2} y(x, t) - c^2 \left(\frac{\partial^2}{\partial x^2} y(x, t) \right) = 0 \quad (1)$$


> $CondFrot := y(0, t) = 0, y(1, t) = 0$

$$CondFrot := y(0, t) = 0, y(1, t) = 0 \quad (2)$$

> $CondIni := y(x, 0) = \frac{\left(\frac{5}{1000}\right)}{\left(\frac{5}{10}\right)} \cdot x \cdot \text{Heaviside}(x) - 2 \cdot \frac{\left(\frac{5}{1000}\right)}{\left(\frac{5}{10}\right)} \cdot \left(x - \frac{5}{10}\right) \cdot \text{Heaviside}\left(x - \frac{5}{10}\right) + \frac{\left(\frac{5}{1000}\right)}{\left(\frac{5}{10}\right)} \cdot (x - 1) \cdot \text{Heaviside}(x - 1)$

$$CondIni := y(x, 0) = \frac{1}{100} x \text{Heaviside}(x) - \frac{1}{50} \left(x - \frac{1}{2}\right) \text{Heaviside}\left(x - \frac{1}{2}\right) + \frac{1}{100} (x - 1) \text{Heaviside}(x - 1) \quad (3)$$

> $\text{plot}(rhs(CondIni), x = -0.2 .. 1.2)$



> $CondIniVel := \text{DerYcero} = 0$

$$CondIniVel := \text{DerYcero} = 0 \quad (4)$$

> $EcuaSep := \text{eval}(\text{subs}(y(x, t) = F(x) \cdot G(t), EDP))$

$$EcuaSep := F(x) \left(\frac{d^2}{dt^2} G(t) \right) - c^2 \left(\frac{d^2}{dx^2} F(x) \right) G(t) = 0 \quad (5)$$

> $EcuaSeparada := \frac{(lhs(EcuaSep) + c \cdot 2 \cdot \text{diff}(F(x), x\$2) \cdot G(t))}{F(x) \cdot G(t)}$

$$= \frac{(rhs(EcuaSep) + c \cdot 2 \cdot \text{diff}(F(x), x\$2) \cdot G(t))}{F(x) \cdot G(t)} \\ EcuaSeparada := \frac{\frac{d^2}{dt^2} G(t)}{G(t)} = \frac{c^2 \left(\frac{d^2}{dx^2} F(x) \right)}{F(x)} \quad (6)$$

> $EcuacionX := \text{rhs}(EcuaSeparada) = \text{alpha}; EcuacionT := \text{lhs}(EcuaSeparada) = \text{alpha}$

$$EcuacionX := \frac{c^2 \left(\frac{d^2}{dx^2} F(x) \right)}{F(x)} = \alpha$$

$$EcuacionT := \frac{\frac{d^2}{dt^2} G(t)}{G(t)} = \alpha \quad (7)$$

> $EcuacionXneg := \text{subs}(\alpha = -n \cdot 2 \cdot c \cdot 2 \cdot \text{Pi} \cdot 2, EcuacionX)$

$$EcuacionXneg := \frac{c^2 \left(\frac{d^2}{dx^2} F(x) \right)}{F(x)} = -n^2 c^2 \pi^2 \quad (8)$$

> $SolucionX := \text{dsolve}(EcuacionXneg)$

$$SolucionX := F(x) = _C1 \sin(\pi n x) + _C2 \cos(\pi n x) \quad (9)$$

> $EcuacionTneg := \text{subs}(\alpha = -n \cdot 2 \cdot c \cdot 2 \cdot \text{Pi} \cdot 2, EcuacionT)$

$$EcuacionTneg := \frac{\frac{d^2}{dt^2} G(t)}{G(t)} = -n^2 c^2 \pi^2 \quad (10)$$

> $SolucionT := \text{dsolve}(EcuacionTneg)$

$$SolucionT := G(t) = _C1 \sin(\pi c n t) + _C2 \cos(\pi c n t) \quad (11)$$

> $SolucionPartX := \text{subs}(_C2 = 0, _C1 = 1, SolucionX)$

$$SolucionPartX := F(x) = \sin(\pi n x) \quad (12)$$

> $SolucionPart := y(x, t) = \text{rhs}(SolucionPartX) \cdot \text{rhs}(SolucionT)$

$$SolucionPart := y(x, t) = \sin(\pi n x) (_C1 \sin(\pi c n t) + _C2 \cos(\pi c n t)) \quad (13)$$

> $SolucionGeneral := y(x, t) = \text{Sum}(\sin(n \cdot \text{Pi} \cdot x) \cdot (b[n] \cos(\text{Pi} \cdot c \cdot n \cdot t) + a[n] \cdot \sin(n \cdot \text{Pi} \cdot c \cdot t))), n = 1 .. \text{infinity}$

$$SolucionGeneral := y(x, t) = \sum_{n=1}^{\infty} \sin(\pi n x) (b_n \cos(\pi c n t) + a_n \sin(\pi c n t)) \quad (14)$$

> $SolucionInicial := \text{eval}(\text{subs}(t = 0, SolucionGeneral))$

$$SolucionInicial := y(x, 0) = \sum_{n=1}^{\infty} \sin(\pi n x) b_n \quad (15)$$

> $b[n] := \frac{1}{\left(\frac{5}{10}\right)} \cdot \text{int}\left(\text{rhs}(CondIni) \cdot \sin\left(\frac{n \cdot \text{Pi} \cdot x}{1}\right), x = 0 .. 1\right)$

$$b_n := \frac{1}{50} \frac{-\sin(\pi n) + 2 \sin\left(\frac{1}{2} \pi n\right)}{\pi^2 n^2} \quad (16)$$

> $a[n] := 0$

$$a_n := 0 \quad (17)$$


> $SolucionGeneral$

$$y(x, t) = \sum_{n=1}^{\infty} \frac{1}{50} \frac{\sin(\pi n x) \left(-\sin(\pi n) + 2 \sin\left(\frac{1}{2} \pi n\right) \right) \cos(\pi c n t)}{\pi^2 n^2} \quad (18)$$

```

> SolucionParticula := y(x, t) =  $\sum_{n=1}^{500}$ 
    $\frac{1}{50} \frac{\sin(\pi n x) \left( -\sin(\pi n) + 2 \sin\left(\frac{1}{2} \pi n\right) \right) \cos(\pi c n t)}{n^2 \pi^2} :$ 
> plot(rhs(subs(c = 1, t = 0, SolucionParticula)), x = 0 .. 1, y = 0 .. 0.007)

```



```

> with(plots) :
> animate(rhs(subs(c = 1, SolucionParticula)), x = 0 .. 1, t = 0 .. 4, frames = 150, view = [0 .. 1, -0.01 .. 0.01])

```

